

**CCDC21 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58824****Specification**

---

**CCDC21 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF
Primary Accession	<a href="#">Q6P2H3</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85639

**CCDC21 Polyclonal Antibody - Additional Information****Gene ID** 64793**Other Names**

Centrosomal protein of 85 kDa, Cep85, Coiled-coil domain-containing protein 21, CEP85, CCDC21

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**CCDC21 Polyclonal Antibody - Protein Information****Name** CEP85 ([HGNC:25309](#))**Synonyms** CCDC21**Function**

Acts as a regulator of centriole duplication through a direct interaction with STIL, a key factor involved in the early steps of centriole formation. The CEP85-STIL protein complex acts as a modulator of PLK4-driven cytoskeletal rearrangements and directional cell motility (PubMed:<a href="http://www.uniprot.org/citations/29712910" target="\_blank">29712910</a>, PubMed:<a href="http://www.uniprot.org/citations/32107292" target="\_blank">32107292</a>). Acts as a negative regulator of NEK2 to maintain the centrosome integrity in interphase. Suppresses centrosome disjunction by inhibiting NEK2 kinase activity (PubMed:<a href="http://www.uniprot.org/citations/26220856" target="\_blank">26220856</a>).

**Cellular Location**

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Nucleus, nucleolus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cell cortex. Note=Localizes to centrosomes and nucleolus in interphase. Upon entry into mitosis, relocates from nucleolus and accumulates at spindle poles

(PubMed:21399614). Associated with the pericentriolar material Localizes to centrosomes at a low level in G1 phase and a slightly increased level in S phase, with gradually elevated levels during G2 phase. The levels at centrosomes further increase at G2/M, reaching a peak at spindle poles at early mitotic stages and remain high until the end of anaphase (PubMed:26220856). Localizes at the leading edge (PubMed:32107292).

### **CCDC21 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **CCDC21 Polyclonal Antibody - Images**