

COMMD4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58840**Specification**

COMMD4 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9H0A8
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21764

COMMD4 Polyclonal Antibody - Additional Information**Gene ID** 54939**Other Names**

COMM domain-containing protein 4, COMMD4

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

COMMD4 Polyclonal Antibody - Protein Information**Name** COMMD4**Function**May modulate activity of cullin-RING E3 ubiquitin ligase (CRL) complexes (PubMed:21778237).
Down-regulates activation of NF- kappa-B.**Cellular Location**

Cytoplasm. Nucleus

Tissue Location


Ubiquitous..

COMMD4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


COMMD4 Polyclonal Antibody - Images



Tissue/cell: rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;


Incubation: Anti-COMMD4 Polyclonal Antibody, Unconjugated(bs-8037R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: humun laryngocarcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-COMMD4 Polyclonal Antibody, Unconjugated(bs-8037R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: human laryngocarcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-COMMD4 Polyclonal Antibody, Unconjugated(bs-8037R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining