

SRRM2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58857**Specification**

SRRM2 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O9UQ35
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	299 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SRRM2
Epitope Specificity	101-230/2752
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus speckle.
SIMILARITY	Belongs to the CWC21 family.
SUBUNIT	Component of the active spliceosome. Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRP70, SNRPA1, SRRM1 and SRRM2. Identified in the spliceosome C complex.
Post-translational modifications	Phosphorylated upon DNA damage, probably by ATM or ATR.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

The SRm160/300 splicing coactivator, which consists of the serine/arginine (SR)-related nuclear matrix protein and a nuclear matrix antigen, functions in splicing by promoting critical interactions between splicing factors bound to pre-mRNA. This splicing pathway involves five core small nuclear ribonucleoprotein particles (snRNPs) and the SR family proteins, which coordinately bind to pre-mRNA splicing enhancer elements, are required for accurate splice site recognition, and regulate alternative splicing patterns. The recognized splicing enhancer elements, known also as exonic enhancer splicing sequences, are short RNA sequences that are capable of activating weak splice sites in adjacent introns and contain specific binding sites for the serine/arginine (SR)-rich splicing factors. SRm160 and 300 antigens contain domains rich in SR motifs, but are distinctly different from the SR factors as they lack an RNA recognition motif and cannot directly induce RNA splicing. These proteins rather function as coactivators that stabilize the splicing complex and mediate the U1 snRNP-splicing pathway.

SRRM2 Polyclonal Antibody - Additional Information

Gene ID 23524**Other Names**

Serine/arginine repetitive matrix protein 2, 300 kDa nuclear matrix antigen, Serine/arginine-rich splicing factor-related nuclear matrix protein of 300 kDa, SR-related nuclear matrix protein of 300 kDa, Ser/Arg-related nuclear matrix protein of 300 kDa, Splicing coactivator subunit SRm300, Tax-responsive enhancer element-binding protein 803, TaxREB803, SRRM2, KIAA0324, SRL300, SRM300

Target/Specificity

Expressed in liver, placenta, and white blood cells.

Dilution

IHC-P ~ N/A
IHC-F ~ N/A
IF ~ 1:50 ~ 200
E ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SRRM2 Polyclonal Antibody - Protein Information

Name SRRM2

Synonyms KIAA0324, SRL300, SRM300

Function

Required for pre-mRNA splicing as component of the spliceosome. As a component of the minor spliceosome, involved in the splicing of U12-type introns in pre-mRNAs (Probable).

Cellular Location

Nucleus. Nucleus speckle

Tissue Location


Expressed in liver, placenta, and white blood cells.

SRRM2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SRRM2 Polyclonal Antibody - Images



Tissue/cell: mouse liver tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-SRRM2 Polyclonal Antibody, Unconjugated(bs-8067R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining