

CCDC34 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58871**Specification**

CCDC34 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O96HJ3
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43225

CCDC34 Polyclonal Antibody - Additional Information**Gene ID** 91057**Other Names**

Coiled-coil domain-containing protein 34, Renal carcinoma antigen NY-REN-41, CCDC34, RAMA3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CCDC34 Polyclonal Antibody - Protein Information**Name** CCDC34**Synonyms** RAMA3**Function**

Involved in spermatogenesis. Has a probable role in anterograde intraflagellar transport which is essential for the formation of sperm flagella.

Cellular Location

Cell projection, cilium, flagellum. Note=Mainly located in the mid-piece of sperm flagella.

Tissue Location


Expressed in sperm..

CCDC34 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC34 Polyclonal Antibody - Images



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CCDC34 Polyclonal Antibody, Unconjugated(bs-8098R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining