

FAM113A Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58918**Specification**

FAM113A Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9H1Q7
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51782

FAM113A Polyclonal Antibody - Additional Information**Gene ID** 64773**Other Names**

PC-esterase domain-containing protein 1A, Protein FAM113A, Sarcoma antigen NY-SAR-23, PCED1A, C20orf81, FAM113A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FAM113A Polyclonal Antibody - Protein Information**Name** PCED1A**Synonyms** C20orf81, FAM113A**FAM113A Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


FAM113A Polyclonal Antibody - Images



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-FAM113A Polyclonal Antibody, Unconjugated(bs-8213R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Human skin tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-FAM113A Polyclonal Antibody, Unconjugated(bs-8213R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining