

FCHSD1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58972**Specification**

FCHSD1 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q86WN1
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76942

FCHSD1 Polyclonal Antibody - Additional Information**Gene ID** 89848**Other Names**

F-BAR and double SH3 domains protein 1, Protein nervous wreck 2, FCHSD1

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FCHSD1 Polyclonal Antibody - Protein Information**Name** FCHSD1**Function**

Promotes actin polymerization mediated by SNX9 and WASL.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q6PFY1}. Perikaryon {ECO:0000250|UniProtKB:Q6PFY1}. Cell projection {ECO:0000250|UniProtKB:Q6PFY1}. Cytoplasmic vesicle {ECO:0000250|UniProtKB:Q6PFY1}. Note=Detected on neuronal cell bodies and cell projections, in part on cytoplasmic vesicles {ECO:0000250|UniProtKB:Q6PFY1}


FCHSD1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FCHSD1 Polyclonal Antibody - Images



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-FCHSD1 Polyclonal Antibody, Unconjugated(bs-8344R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining