

Ube2H Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58978**Specification****Ube2H Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P62256
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20655

Ube2H Polyclonal Antibody - Additional Information**Gene ID** 7328**Other Names**

Ubiquitin-conjugating enzyme E2 H, 2.3.2.23, (E3-independent) E2 ubiquitin-conjugating enzyme H, 2.3.2.24, E2 ubiquitin-conjugating enzyme H, UbcH2, Ubiquitin carrier protein H, Ubiquitin-conjugating enzyme E2-20K, Ubiquitin-protein ligase H, UBE2H

Dilution

WB~1:1000<br \>IHC-P~N/A<br \>IHC-F~N/A<br \>IF~1:50~200<br \>E~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Ube2H Polyclonal Antibody - Protein Information**Name** UBE2H**Function**

Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins (PubMed:17588522, PubMed:20061386, PubMed:8132613). E2 ubiquitin conjugating enzyme that transfers ubiquitin to MAEA, a core component of the CTLH E3 ubiquitin- protein ligase complex (PubMed:29911972). In vitro catalyzes 'Lys- 11'- and 'Lys-48'-linked polyubiquitination (PubMed:20061386). Capable, in vitro, to ubiquitinate histone H2A (PubMed:8132613).

Ube2H Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ube2H Polyclonal Antibody - Images