

NCE2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58982**Specification****NCE2 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q969M7
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NCE2
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the ubiquitin-conjugating enzyme family. UBE2F subfamily.
SUBUNIT	Interacts with UBA3 and RBX2.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Accepts the ubiquitin-like protein NEDD8 from the UBA3-NAE1 E1 complex and catalyzes its covalent attachment to other proteins. The specific interaction with the E3 ubiquitin ligase RBX2, but not RBX1, suggests that the RBX2-UBE2F complex neddylates specific target proteins, such as CUL5.

NCE2 Polyclonal Antibody - Additional Information**Gene ID** 140739**Other Names**

NEDD8-conjugating enzyme UBE2F, 2.3.2.32, NEDD8 carrier protein UBE2F, NEDD8 protein ligase UBE2F, NEDD8-conjugating enzyme 2, RING-type E3 NEDD8 transferase UBE2F, Ubiquitin-conjugating enzyme E2 F, UBE2F, NCE2

Target/Specificity

Widely expressed.

Dilution

dilution_WB WB~~1:1000
dilution_IHC-P IHC-P~~N/A
dilution_IHC-F IHC-F~~N/A

=dilution_IF">IF~1:50~200<br \>E~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NCE2 Polyclonal Antibody - Protein Information

Name UBE2F

Synonyms NCE2

Function

Accepts the ubiquitin-like protein NEDD8 from the UBA3-NAE1 E1 complex and catalyzes its covalent attachment to other proteins (PubMed:19250909, PubMed:23201271). Together with the E3 ubiquitin ligase RNF7/RBX2, specifically neddylates cullin-5 (CUL5) (PubMed:19250909, PubMed:23201271, PubMed:23300442). Does not neddylate CUL1, CUL2, CUL3, CUL4A or CUL4B (PubMed:19250909, PubMed:23201271). Mediates neddylation of the CUL9-RBX1 complex (PubMed:38605244).

Tissue Location

Widely expressed (at protein level).

NCE2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCE2 Polyclonal Antibody - Images