

OLIG1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59034

Specification

OLIG1 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, E

Primary Accession Q8TAK6

Reactivity
Host
Clonality
Calculated MW
Physical State

Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
28 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human OLIG1

Epitope Specificity 121-220/271

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus.

SIMILARITY Contains 1 bHLH (basic helix-loop-helix)

domain.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

The oligodendrocyte lineage-specific basic helix-loop-helix (OLIG) family of transcription factors include OLIG1-OLIG3, which differ in tissue expression. OLIG1 and OLIG2 are specifically expressed in nervous tissue as gene regulators of oligodendrogenesis. OLIG2 is more widely expressed in embryonic brain than OLIG1, while OLIG3 is primarily expressed in non-neural tissues. OLIG1 and OLIG2 interact with the Nkx-2.2 homeodomain protein, which is responsible for directing ventral neuronal patterning in response to graded Sonic hedgehog signaling in the embryonic neural tube. These interactions between OLIG proteins and Nkx-2.2 appear to promote the formation of alternate cell types by inhibiting V3 interneuron development. OLIG1 and OLIG2 are abundantly expressed in oligodendroglioma and nearly absent in astrocytomas. Therefore, OLIG proteins are candidates for molecular markers of human glial brain tumors, which are the most common primary malignancies of the human brain.

OLIG1 Polyclonal Antibody - Additional Information

Gene ID 116448

Other Names

Oligodendrocyte transcription factor 1, Oligo1, Class B basic helix-loop-helix protein 6, bHLHb6, Class E basic helix-loop-helix protein 21, bHLHe21, OLIG1, BHLHB6, BHLHE21



Target/Specificity

Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas is highly variable.

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

OLIG1 Polyclonal Antibody - Protein Information

Name OLIG1

Synonyms BHLHB6, BHLHE21

Function

Promotes formation and maturation of oligodendrocytes, especially within the brain. Cooperates with OLIG2 to establish the pMN domain of the embryonic neural tube (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.

Tissue Location

Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas is highly variable.

OLIG1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

OLIG1 Polyclonal Antibody - Images