

SARM1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59096

Specification

SARM1 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, E

Primary Accession 06SZW1 Rat, Pig, Dog, Bovine Reactivity

Host Rabbit Clonality **Polyclonal** Calculated MW 80 KDa **Physical State** Liquid

Immunogen KLH conjugated synthetic peptide derived

from human SARM1

321-420/724 **Epitope Specificity**

Isotype laG **Purity**

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm.

SIMILARITY

Contains 2 SAM (sterile alpha motif) domains. Contains 1 TIR domain.

SUBUNIT Interacts with TICAM1/TRIF and thereby interferes with TICAM1/TRIF function. This product as supplied is intended for Important Note research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

SARM, also known as SAMD2, SARM1 or KIAA0524, is a 724 amino acid protein that localizes to the cytoplasm and contains one TIR domain and two sterile alpha motif (SAM) domains. Expressed predominately in liver and kidney and present at lower levels in placenta, SARM interacts with TICAM-1 and, via this interaction, blocks the transcriptional activation activity of TICAM-1 and functions as a negative regulator of Toll-like receptor signaling. Additionally, SARM is thought to be involved in innate immune responses and may also play a role in the negative regulation of NFxB activation. SARM exists as two alternatively spliced isoforms that are encoded by a gene which maps to human chromosome 17.

SARM1 Polyclonal Antibody - Additional Information

Gene ID 23098

Other Names

NAD(+) hydrolase SARM1, NADase SARM1, hSARM1, 3.2.2.6, NADP(+) hydrolase SARM1, 3.2.2.-, Sterile alpha and Armadillo repeat protein, Sterile alpha and TIR motif-containing protein 1 {ECO:0000303|PubMed:18089857, ECO:0000303|Ref.2}, Sterile alpha motif domain-containing protein 2, MyD88-5, SAM domain-containing protein 2 {ECO:0000312|HGNC:HGNC:17074}, Tir-1



homolog, HsTIR, SARM1

Target/Specificity

Predominantly expressed in kidney and liver. Expressed at lower level in placenta.

Dilution

```
<span class ="dilution_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution_IF">IF~~1:50~200</span><br \> <span class ="dilution_E">E~~N/A</span>
```

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

SARM1 Polyclonal Antibody - Protein Information

Name SARM1

Function

NAD(+) hydrolase, which plays a key role in axonal degeneration following injury by regulating NAD(+) metabolism (PubMed:25908823, PubMed:27671644, PubMed:28334607). Acts as a negative regulator of MYD88- and TRIF-dependent toll-like receptor signaling pathway by promoting Wallerian degeneration, an injury-induced form of programmed subcellular death which involves degeneration of an axon distal to the injury site $(PubMed:<a\ href="http://www.uniprot.org/citations/15123841" \ target="_blank">15123841^, PubMed:16964262,$ PubMed:20306472, PubMed:25908823). Wallerian degeneration is triggered by NAD(+) depletion: in response to injury, SARM1 is activated and catalyzes cleavage of NAD(+) into ADP-D-ribose (ADPR), cyclic ADPR (cADPR) and nicotinamide; NAD(+) cleavage promoting cytoskeletal degradation and axon destruction (PubMed:25908823, PubMed:28334607, PubMed:30333228, PubMed: 31128467, PubMed:31439792, PubMed:31439793, PubMed: 32049506, PubMed: 32828421, $PubMed: 33053563).$ Also able to hydrolyze NADP(+), but not other NAD(+)-related molecules (PubMed:29395922). Can activate neuronal cell death in response to stress (PubMed:20306472). Regulates dendritic arborization through the MAPK4-JNK pathway (By similarity). Involved in innate immune response: inhibits both TICAM1/TRIF- and MYD88-dependent activation of JUN/AP-1, TRIF-dependent activation of NF-kappa-B and IRF3, and the phosphorylation of MAPK14/p38 (PubMed:16964262).

Cellular Location

Cytoplasm. Cell projection, axon {ECO:0000250|UniProtKB:Q6PDS3}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q6PDS3}. Synapse {ECO:0000250|UniProtKB:Q6PDS3}. Mitochondrion Note=Associated with microtubules. {ECO:0000250|UniProtKB:Q6PDS3}



Tissue Location


Predominantly expressed in brain, kidney and liver. Expressed at lower level in placenta.

SARM1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SARM1 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Human liver tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SARM1) Polyclonal Antibody, Unconjugated (bs-9001R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.