

AHNAK2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59139**Specification**

AHNAK2 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q8IVF2
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	616629

AHNAK2 Polyclonal Antibody - Additional Information**Gene ID** 113146**Other Names**

Protein AHNAK2, AHNAK2, C14orf78, KIAA2019

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

AHNAK2 Polyclonal Antibody - Protein Information**Name** AHNAK2**Synonyms** C14orf78, KIAA2019**Cellular Location**

Nucleus.


AHNAK2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

AHNAK2 Polyclonal Antibody - Images



Sample:


Cerebrum (Mouse) Lysate at 40 ug

Primary: Anti- AHNAK2 (bs-9094R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 616 kD

Observed band size: 70 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (AHNAK2) Polyclonal Antibody, Unconjugated (bs-9094R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.