

ZNRF1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59157

Specification

ZNRF1 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, E

Primary Accession Q8ND25

Reactivity Rat, Dog, Bovine Host Rabbit Clonality **Polyclonal** Calculated MW **24 KDa** Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human ZNRF1

121-227/227 **Epitope Specificity**

laG Isotype **Purity**

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol. SUBCELLULAR LOCATION **Endosome.** Lysosome. Membrane.

SIMILARITY Contains 1 RING-type zinc finger.

Interacts with AKT1, GLUL and TUBB2A (By **SUBUNIT**

similarity).

This product as supplied is intended for Important Note

research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. ZNRF1 (zinc and ring finger 1), also known as NIN283, is a 227 amino acid protein that contains one RING-type zinc finger and localizes to the lysosome and the endosome, as well as to cytoplasmic vesicles and the peripheral membrane. Expressed primarily in nervous system tissue, but also present in testis and thymus, ZNRF1 functions as an E3 ubiquitin-protein ligase that is thought to play a role in the establishment and maintenance of neuronal plasticity. Multiple isoforms of ZNRF1 exist due to alternative splicing events.

ZNRF1 Polyclonal Antibody - Additional Information

Gene ID 84937

Other Names

E3 ubiquitin-protein ligase ZNRF1, 2.3.2.27, Nerve injury-induced gene 283 protein, RING-type E3 ubiquitin transferase ZNRF1, Zinc/RING finger protein 1, ZNRF1, NIN283



Target/Specificity

Expressed primarily in the nervous system, with expression higher in developing brain relative to adult. Expressed at low levels in testis and thymus.

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

ZNRF1 Polyclonal Antibody - Protein Information

Name ZNRF1

Synonyms NIN283

Function

E3 ubiquitin-protein ligase that plays a role in different processes including cell differentiation, receptor recycling or regulation of inflammation (PubMed:28593998, PubMed:33996800, PubMed:37158982). Mediates the ubiquitination of AKT1 and GLUL, thereby playing a role in neuron cells differentiation. Plays a role in the establishment and maintenance of neuronal transmission and plasticity. Regulates Schwann cells differentiation by mediating ubiquitination of GLUL. Promotes neurodegeneration by mediating 'Lys-48'-linked polyubiquitination and subsequent degradation of AKT1 in axons: degradation of AKT1 prevents AKT1-mediated phosphorylation of GSK3B, leading to GSK3B activation and phosphorylation of DPYSL2/CRMP2 followed by destabilization of microtubule assembly in axons. Ubiquitinates the Na(+)/K(+) ATPase alpha-1 subunit/ATP1A1 and thereby influences its endocytosis and/or degradation (PubMed:22797923). Controls ligand-induced EGFR signaling via mediating receptor ubiquitination and recruitment of the ESCRT machinery (PubMed:<a href="http://www.uniprot.org/citations/33996800"

target="_blank">33996800). Acts as a negative feedback mechanism controlling TLR3 trafficking by mediating TLR3 'Lys-63'-linked polyubiquitination to reduce type I IFN production (PubMed:37158982). Modulates inflammation by promoting caveolin-1/CAV1 ubiquitination and degradation to regulate TLR4-activated immune response (PubMed:28593998).


Cellular Location

Endosome. Lysosome. Membrane; Peripheral membrane protein. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane; Peripheral membrane protein Note=Associated with synaptic vesicle membranes in neurons

Tissue Location

Expressed primarily in the nervous system, with expression higher in developing brain relative to adult. Expressed at low levels in testis and thymus.

ZNRF1 Polyclonal Antibody - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZNRF1 Polyclonal Antibody - Images