

UBR2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59159

Specification

UBR2 Polyclonal Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, IHC-P, IHC-F, IF, E
O8IWV8
Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
200538

UBR2 Polyclonal Antibody - Additional Information

Gene ID 23304

Other Names

E3 ubiquitin-protein ligase UBR2, 2.3.2.27, N-recognin-2, RING-type E3 ubiquitin transferase UBR2, Ubiquitin-protein ligase E3-alpha-2, Ubiquitin-protein ligase E3-alpha-II, UBR2, C6orf133, KIAA0349

Dilution

```
<span class ="dilution_WB">WB~~1:1000</span><br \> <span class
="dilution_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution_IF">IF~~1:50~200</span><br \> <span class ="dilution_E">E~~N/A</span>
```

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

UBR2 Polyclonal Antibody - Protein Information

Name UBR2

Synonyms C6orf133, KIAA0349

Function

E3 ubiquitin-protein ligase which is a component of the N-end rule pathway (PubMed:15548684, PubMed:20835242, PubMed:28392261). Recognizes and binds to proteins bearing specific N-terminal residues (N-degrons) that are destabilizing according to the N-end rule, leading to their ubiquitination and subsequent degradation (PubMed:20835242, PubMed:20835242, Recognizes both type-1 and type-2 N-degrons, containing positively charged amino acids (Arg, Lys and His) and bulky and hydrophobic amino acids, respectively (PubMed:<a



 $href="http://www.uniprot.org/citations/20835242" target="_blank">20835242, PubMed:28392261). Does not ubiquitinate proteins that are acetylated at the N-terminus (PubMed:<a$

href="http://www.uniprot.org/citations/20835242" target="_blank">20835242). In contrast, it strongly binds methylated N-degrons (PubMed:28392261). Plays a critical role in chromatin inactivation and chromosome-wide transcriptional silencing during meiosis via ubiquitination of histone H2A (By similarity). Binds leucine and is a negative regulator of the leucine-mTOR signaling pathway, thereby controlling cell growth (PubMed:20298436). Required for spermatogenesis, promotes, with Tex19.1, SPO11-dependent recombination foci to accumulate and drive robust homologous chromosome synapsis (By similarity). Polyubiquitinates LINE-1 retrotransposon encoded, LIRE1, which induces degradation, inhibiting LINE-1 retrotransposon mobilization (By similarity). Catalyzes ubiquitination and degradation of the N-terminal part of NLRP1 following NLRP1 activation by pathogens and other damage-associated signals: ubiquitination promotes degradation of the N-terminal part and subsequent release of the cleaved C-terminal part of NLRP1, which polymerizes and forms the NLRP1 inflammasome followed by host cell pyroptosis (By similarity). Plays a role in T-cell receptor signaling by inducing 'Lys-63'-linked ubiquitination of lymphocyte cell-specific kinase LCK (PubMed:38225265). This activity is regulated by DUSP22, which induces 'Lys-48'-linked ubiquitination of UBR2, leading to its proteasomal degradation by SCF E3 ubiquitin-protein ligase complex (PubMed:38225265).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q6WKZ8}. Chromosome {ECO:0000250|UniProtKB:Q6WKZ8}. Note=Associated with chromatin during meiosis. {ECO:0000250|UniProtKB:Q6WKZ8}

Tissue Location

Broadly expressed, with highest levels in skeletal muscle, kidney and pancreas. Present in acinar cells of the pancreas (at protein level).

UBR2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

UBR2 Polyclonal Antibody - Images