

VGLL3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59174**Specification**

VGLL3 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	A8MV65
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36009

VGLL3 Polyclonal Antibody - Additional Information**Gene ID** 389136**Other Names**Transcription cofactor vestigial-like protein 3, Vgl-3, VGLL3
{ECO:0000312|Ensembl:ENSP00000381436}**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

VGLL3 Polyclonal Antibody - Protein Information**Name** VGLL3**Function**

May act as a specific coactivator for the mammalian TEFs.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q8N8G2}.

Tissue Location

Enriched in placenta.


VGLL3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VGLL3 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (VGLL3) Polyclonal Antibody, Unconjugated (bs-9184R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.