

**Cystatin D Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP59183****Specification**

---

**Cystatin D Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC-P, IHC-F, IF, E    |
| Primary Accession | <a href="#">P28325</a> |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 16080                  |

**Cystatin D Polyclonal Antibody - Additional Information****Gene ID** 1473**Other Names**

Cystatin-D, Cystatin-5, CST5

**Dilution**

IHC-P~N/A  
IHC-F~N/A  
IF~1:50~200  
E~N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Cystatin D Polyclonal Antibody - Protein Information****Name** CST5**Function**

Cysteine proteinase inhibitor that possibly plays a protective role against proteinases present in the oral cavity. The order of preference for inhibition is cathepsin S > cathepsin H > cathepsin L > cathepsin B.

**Cellular Location**

Secreted.

**Tissue Location**

Expressed in submandibular and sublingual saliva but not in parotid saliva (at protein level).  
Expressed in parotid gland but not in seminal vesicle, prostate, epididymis, testis, ovary, placenta, thyroid, gastric corpus, small intestine, liver, or gall bladder tissue.

## **Cystatin D Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Cystatin D Polyclonal Antibody - Images**