

RNF156 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59187**Specification**

RNF156 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O60291
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60753

RNF156 Polyclonal Antibody - Additional Information**Gene ID** 23295**Other Names**

E3 ubiquitin-protein ligase MGRN1, 2.3.2.27, Mahogunin RING finger protein 1, RING finger protein 156, RING-type E3 ubiquitin transferase MGRN1, MGRN1, KIAA0544, RNF156

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RNF156 Polyclonal Antibody - Protein Information**Name** MGRN1**Synonyms** KIAA0544, RNF156**Function**

E3 ubiquitin-protein ligase. Mediates monoubiquitination at multiple sites of TSG101 in the presence of UBE2D1, but not of UBE2G1, nor UBE2H. Plays a role in the regulation of endosome-to-lysosome trafficking. Impairs MC1R- and MC4R-signaling by competing with GNAS-binding to MCRs and inhibiting agonist-induced cAMP production. Does not inhibit ADRB2-signaling. Does not promote MC1R ubiquitination. Acts also as a negative regulator of hedgehog signaling (By similarity).

Cellular Location


Early endosome. Note=The endosomal localization is dependent on the interaction with TSG101 [Isoform 2]: Cytoplasm, cytosol. Nucleus. Note=Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. Excluded from nucleoli [Isoform 4]: Cytoplasm, cytosol. Cell membrane.

RNF156 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RNF156 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (mouse stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF156) Polyclonal Antibody, Unconjugated (bs-9229R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining



Paraformaldehyde-fixed, paraffin embedded (Human hepatocellular carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF156) Polyclonal Antibody, Unconjugated (bs-9229R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining