

### **NDE1 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59218

# **Specification**

### **NDE1 Polyclonal Antibody - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

IHC-P, IHC-F, IF, E

O9NXR1

Rat, Pig, Dog

Rabbit

Polyclonal

37721

# **NDE1 Polyclonal Antibody - Additional Information**

Gene ID 54820

#### **Other Names**

Nuclear distribution protein nudE homolog 1, NudE, NDE1, NUDE

#### **Dilution**

```
<span class ="dilution_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution_IF">IF~~1:50~200</span><br \> <span class = "dilution_E">E~~N/A</span>
```

### Storage

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.


# **NDE1 Polyclonal Antibody - Protein Information**

Name NDE1 (HGNC:17619)

**Synonyms NUDE** 

#### **Function**

Required for centrosome duplication and formation and function of the mitotic spindle. Essential for the development of the cerebral cortex. May regulate the production of neurons by controlling the orientation of the mitotic spindle during division of cortical neuronal progenitors of the proliferative ventricular zone of the brain. Orientation of the division plane perpendicular to the layers of the cortex gives rise to two proliferative neuronal progenitors whereas parallel orientation of the division plane yields one proliferative neuronal progenitor and a postmitotic neuron. A premature shift towards a neuronal fate within the progenitor population may result in an overall reduction in the final number of neurons and an increase in the number of neurons in the deeper layers of the cortex. Acts as a RAB9A/B effector that tethers RAB9-associated late endosomes to the dynein motor for their retrograde transport to the trans-Golgi network (PubMed:<a href="http://www.uniprot.org/citations/34793709" target="\_blank">>34793709</a>/a>).





#### **Cellular Location**

Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, spindle. Cleavage furrow Cytoplasmic vesicle membrane Note=Localizes to the interphase and S phase centrosome. During mitosis, partially associated with the mitotic spindle. Concentrates at the plus ends of microtubules coincident with kinetochores in metaphase and anaphase in a CENPF-dependent manner. Also localizes to the cleavage furrow during cytokinesis. manner. Also localizes to the cleavage furrow during cytokinesis. Colocalizes with RAB9A to membrane vesicles (PubMed:34793709).

# **Tissue Location**

Expressed in the neuroepithelium throughout the developing brain, including the cerebral cortex and cerebellum

# **NDE1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

NDE1 Polyclonal Antibody - Images