

MARCH2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59227

Specification

MARCH2 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, E

Primary Accession
Reactivity
Rat
Host
Clonality
Calculated MW
Rat
Rabbit
Polyclonal
27 KDa

Physical State
Liquid
Immunogen
KLH conjugated synthetic peptide derived

from human MARCH2

Epitope Specificity 151-246/246

Isotype IgG

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Endoplasmic reticulum membrane.

Lysosome membrane. Endosome

membrane.

SIMILARITY Contains 1 RING-CH-type zinc finger.
SUBUNIT Interacts with STX6 (By similarity).

Interacts with MARCH3.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). MARCH2 (membrane-associated ring finger (C3HC4) 2), also known as RNF172 or HSPC240, is a 246 amino acid multi-pass membrane protein that localizes to the endoplasmic reticulum and contains one RING-CH-type zinc finger. Expressed in a variety of tissues, MARCH2 functions as an E3 ubiquitin-protein ligase that is thought to mediate the ubiquitination and subsequent degradation of CD71 and B7-2 and may be involved in endosomal protein trafficking.

MARCH2 Polyclonal Antibody - Additional Information

Gene ID 51257

Other Names

E3 ubiquitin-protein ligase MARCHF2, 2.3.2.27, Membrane-associated RING finger protein 2, Membrane-associated RING-CH protein II, MARCH-II, RING finger protein 172, RING-type E3



ubiquitin transferase MARCHF2, MARCHF2 (HGNC:28038), MARCH2, RNF172

Target/Specificity Broadly expressed.

Dilution

WB~~1:1000<br \> <span class
="dilution_IHC-P">IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

MARCH2 Polyclonal Antibody - Protein Information

Name MARCHF2 (HGNC:28038)

Synonyms MARCH2, RNF172

Function

E3 ubiquitin-protein ligase that may mediate ubiquitination of TFRC and CD86, and promote their subsequent endocytosis and sorting to lysosomes via multivesicular bodies. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates (PubMed:14722266, PubMed:16428329). Together with GOPC/CAL mediates the ubiquitination and lysosomal degradation of CFTR (PubMed:23818989). Ubiquitinates and therefore mediates the degradation of DLG1 (PubMed:17980554). Regulates the intracellular trafficking and secretion of alpha1-antitrypsin/SERPINA1 and HP/haptoglobin via ubiquitination and degradation of the cargo receptor ERGIC3 (PubMed:31142615). Negatively regulates the antiviral and antibacterial immune response by repression of the NF-kB and type 1 IFN signaling pathways, via MARCHF2-mediated K48-linked polyubiquitination of IKBKG/NEMO, resulting in its proteasomal degradation (PubMed:32935379). May be involved in endosomal trafficking through interaction with STX6 (PubMed:15689499).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Lysosome membrane; Multi-pass membrane protein. Endosome membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasm. Cell membrane; Multi-pass membrane protein

Tissue Location

Broadly expressed..



Tel: 858.875.1900 Fax: 858.875.1999



MARCH2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

MARCH2 Polyclonal Antibody - Images