

PSMD6 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59235

Specification

PSMD6 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, E

Primary Accession <u>Q15008</u>

Reactivity
Host
Clonality
Calculated MW
Physical State
Rat, Pig, Bovine
Rabbit
Polyclonal
43 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human PSMD6

Epitope Specificity 121-230/389

Isotype IgG
Purity

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Proteasome complex.

SIMILARITY Belongs to the proteasome subunit S10

family.Contains 1 PCI domain.

SUBUNIT

Important Note

Component of the PA700 complex.

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. Proteasome 26S S10 is a non-ATPase subunit of the 19S regulator. It acts as a regulatory subunit of the 26S proteasome which is involved in the ATP-dependent degradation of ubiquitinated proteins. Two transcripts encoding different isoforms have been described. Pseudogenes have been identified on chromosomes 3 and 20.

PSMD6 Polyclonal Antibody - Additional Information

Gene ID 9861

Other Names

26S proteasome non-ATPase regulatory subunit 6, 26S proteasome regulatory subunit RPN7, 26S



proteasome regulatory subunit S10, Breast cancer-associated protein SGA-113M, Phosphonoformate immuno-associated protein 4, Proteasome regulatory particle subunit p44S10, p42A, PSMD6, KIAA0107, PFAAP4

Target/Specificity

Overexpressed in hepatocellular carcinomas.

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

PSMD6 Polyclonal Antibody - Protein Information

Name PSMD6

Synonyms KIAA0107, PFAAP4

Function

Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair.

PSMD6 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

PSMD6 Polyclonal Antibody - Images