

CARKD Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59245

Specification

CARKD Polyclonal Antibody - Product Information

Application Primary Accession

Reactivity
Host
Clonality
Calculated MW
Physical State
Immunogen

Epitope Specificity

Isotype Purity

affinity purified by Protein A

WB, IHC-P, IHC-F, IF, E

Q8IW45 Rat Rabbit Polyclonal 37 KDa Liquid

KLH conjugated synthetic peptide derived

from human CARKD

251-347/347

IqG

Buffer

SUBCELLULAR LOCATION SIMILARITY Important Note 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.
Mitochondrion (By similarity).
Belongs to the nnrD/CARKD family.
This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

CARKD is a 347 amino acid protein that belongs to the YjeF family and exists as three alternatively spliced isoforms. Containing one YjeF C-terminal domain, CARKD is encoded by a gene that maps to human chromosome 13q34. Chromosome 13 contains around 114 million base pairs and 400 genes. Key tumor suppressor genes on chromosome 13 include the breast cancer susceptibility gene, BRCA2, and the RB1 (retinoblastoma) gene. RB1 encodes a crucial tumor suppressor protein which, when defective, leads to malignant growth in the retina and has been implicated in a variety of other cancers. The gene SLITRK1, which is associated with Tourette syndrome, is on chromosome 13. As with most chromosomes, polysomy of part or all of chromosome 13 is deleterious to development and decreases the odds of survival. Trisomy 13, also known as Patau syndrome, is quite deadly and the few who survive past one year suffer from permanent neurologic defects, difficulty eating and vulnerability to serious respiratory infections.

CARKD Polyclonal Antibody - Additional Information

Gene ID 55739

Other Names

 $ATP-dependent (S)-NAD(P)H-hydrate dehydratase \\ \{ECO:0000255|HAMAP-Rule:MF_03157\}, \\ 4.2.1.93 \\ \{ECO:0000255|HAMAP-Rule:MF_03157, ECO:0000269|PubMed:30576410\}, \\ ATP-dependent NAD(P)HX dehydratase \\ \{ECO:0000255|HAMAP-Rule:MF_03157\}, Carbohydrate kinase domain-containing protein \\ \{ECO:0000255|HAMAP-Rule:MF_03157\}, NAD(P)HX dehydratase \\ \{ECO:0000255|HAMAP-Rul$



{ECO:0000312|HGNC:HGNC:25576}, NAXD (HGNC:25576)

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

CARKD Polyclonal Antibody - Protein Information

Name NAXD (HGNC:25576)

Function

Catalyzes the dehydration of the S-form of NAD(P)HX at the expense of ATP, which is converted to ADP. Together with NAD(P)HX epimerase, which catalyzes the epimerization of the S- and R-forms, the enzyme allows the repair of both epimers of NAD(P)HX, a damaged form of NAD(P)H that is a result of enzymatic or heat-dependent hydration.

Cellular Location

Mitochondrion {ECO:0000255|HAMAP-Rule:MF_03157}.


CARKD Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CARKD Polyclonal Antibody - Images





Sample: MCF-7 Cell(Human)Lysate at 30 ug Primary: Anti-CARKD (bs-9402R)at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 37kD Observed band size: 45kD