

KChIP2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59257**Specification****KChIP2 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	O9NS61
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30907

KChIP2 Polyclonal Antibody - Additional Information**Gene ID** 30819**Other Names**

Kv channel-interacting protein 2, KChIP2, A-type potassium channel modulatory protein 2, Cardiac voltage-gated potassium channel modulatory subunit, Potassium channel-interacting protein 2, KCNIP2, KCHIP2

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KChIP2 Polyclonal Antibody - Protein Information**Name** KCNIP2 ([HGNC:15522](#))**Function**

Regulatory subunit of Kv4/D (Shal)-type voltage-gated rapidly inactivating A-type potassium channels (PubMed:10676964, PubMed:11287421, PubMed:11684073, PubMed:12297301, PubMed:14623880, PubMed:34997220). Modulates channel density, inactivation kinetics and rate of

recovery from inactivation in a calcium-dependent and isoform- specific manner (PubMed:10676964, PubMed:11287421, PubMed:11684073, PubMed:12297301, PubMed:14623880, PubMed:34997220). Involved in KCND2 and KCND3 trafficking to the cell surface (PubMed:12829703). May be required for the expression of I(To) currents in the heart (By similarity).

Cellular Location

[Isoform 1]: Cell membrane {ECO:0000250|UniProtKB:Q9JM59}; Lipid-anchor {ECO:0000250|UniProtKB:Q9JM59}. Note=Detected on lipid rafts (By similarity). {ECO:0000250|UniProtKB:Q9JM59} [Isoform 6]: Cell membrane {ECO:0000250|UniProtKB:Q9JM59}; Lipid-anchor {ECO:0000250|UniProtKB:Q9JM59}

Tissue Location

Expressed in brain. Colocalizes with KCND2 in excitatory neurons including cortical and hippocampal CA1 pyramidal cells. Isoform 3 is expressed in heart and in umbilical vein endothelial cells. Not expressed in fetal heart

KChIP2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KChIP2 Polyclonal Antibody - Images