

KRCC1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59291**Specification**

KRCC1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9NPI7
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30980

KRCC1 Polyclonal Antibody - Additional Information**Gene ID** 51315**Other Names**

Lysine-rich coiled-coil protein 1, Cryptogenic hepatitis-binding protein 2, KRCC1, CHBP2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KRCC1 Polyclonal Antibody - Protein Information**Name** KRCC1**Synonyms** CHBP2**KRCC1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KRCC1 Polyclonal Antibody - Images



Sample:

Kidney (Mouse) Lysate at 40 ug


Cerebrum (Mouse) Lysate at 40 ug

Primary: Anti-KRCC1 (bs-9560R) at 1/300 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 31 kD


Observed band size: 31 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse Brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KRCC1) Polyclonal Antibody, Unconjugated (bs9560R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse Kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KRCC1) Polyclonal Antibody, Unconjugated (bs9560R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KRCC1) Polyclonal Antibody, Unconjugated (bs-9560R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.