

C17orf74 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59323**Specification**

C17orf74 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	Q0P670
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57131

C17orf74 Polyclonal Antibody - Additional Information**Gene ID** 201243**Other Names**

Uncharacterized protein SPEM2, SPEM2 (HGNC:27315), C17orf74

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C17orf74 Polyclonal Antibody - Protein Information**Name** SPEM2 ([HGNC:27315](#))**Synonyms** C17orf74**Cellular Location**

Membrane; Single-pass membrane protein


C17orf74 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


C17orf74 Polyclonal Antibody - Images



Tissue/cell: Rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;


Incubation: Anti-C17orf74 Polyclonal Antibody, Unconjugated(bs-9645R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-C17orf74 Polyclonal Antibody, Unconjugated(bs-9645R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample: A549 Cell (Human) Lysate at 40 ug
Primary: Anti-C17orf74 (bs-9645R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 57 kD
Observed band size: 63 kD