

## C17orf75 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59324

### **Specification**

# C17orf75 Polyclonal Antibody - Product Information

Application

Primary Accession
Reactivity
Rat, Dog
Host
Clonality
Calculated MW
Physical State

Q9HASO
Rat, Dog
Rabbit
Polyclonal
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human C17orf75

WB, IHC-P, IHC-F, IF, ICC, E

201-300/396

IgG

**Epitope Specificity** 

Isotype **Purity** 

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

## **Background Descriptions**

C17orf75 is a 396 amino acid protein that is encoded by a gene mapping to human chromosome 17. Chromosome 17 makes up over 2.5% of the human genome with about 81 million bases encoding over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes. Chromosome 17 is also linked to neurofibromatosis, a condition characterized by neural and epidermal lesions, and dysregulated Schwann cell growth. Alexander disease, Birt-Hogg-Dube syndrome and Canavan disease are also associated with chromosome 17.

# C17orf75 Polyclonal Antibody - Additional Information

**Gene ID 64149** 

Other Names

Protein Njmu-R1, C17orf75

### Target/Specificity

The molecular function and cellular component of this protein are unknown. It may have a role in spermatogenesis, is highly expressed in testis and is also expressed in fetal testis.



#### **Dilution**

<span class ="dilution\_WB">WB~~1:1000</span><br \><span class
="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class
="dilution\_IF">IF~~1:50~200</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class ="dilution\_ICC">ICC~~N/A</span>

### **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## C17orf75 Polyclonal Antibody - Protein Information

### Name C17orf75

### **Function**

As component of the WDR11 complex acts together with TBC1D23 to facilitate the golgin-mediated capture of vesicles generated using AP-1 (PubMed:<a href="http://www.uniprot.org/citations/29426865" target="\_blank">29426865</a>). May have a role in spermatogenesis.

#### **Cellular Location**

Golgi apparatus, trans-Golgi network. Cytoplasmic vesicle

### **Tissue Location**

Highly expressed in testis and also expressed in fetal testis


# C17orf75 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## C17orf75 Polyclonal Antibody - Images





Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C17orf75) Polyclonal Antibody, Unconjugated (bs-9646R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



# Sample:

U2os Cell (Human) Lysate at 30 ug

Primary: Anti- C17orf75 (bs-9646R)at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 45kD Observed band size: 45kD