

C20orf11 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59344**Specification**

C20orf11 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	O9NWU2
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26749

C20orf11 Polyclonal Antibody - Additional Information**Gene ID** 54994**Other Names**

Glucose-induced degradation protein 8 homolog, Two hybrid-associated protein 1 with RanBPM, Twa1, GID8

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C20orf11 Polyclonal Antibody - Protein Information**Name** GID8**Function**

Core component of the CTLH E3 ubiquitin-protein ligase complex that selectively accepts ubiquitin from UBE2H and mediates ubiquitination and subsequent proteasomal degradation of the transcription factor HBP1 (PubMed:29911972). Acts as a positive regulator of Wnt signaling pathway by promoting beta-catenin (CTNNB1) nuclear accumulation (PubMed:28829046).

Cellular Location

Cytoplasm. Nucleus. Note=Localizes in the cytoplasm in the absence of Wnt stimulation and in the nucleus in the presence of Wnt stimulation.

Tissue Location

Up-regulated in colorectal cancer tissues (at protein level).

C20orf11 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C20orf11 Polyclonal Antibody - Images