

Kazrin Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59356

Specification

Kazrin Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q674X7
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	86 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Kazrin
Epitope Specificity	451-550/775
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm
SIMILARITY	Belongs to the kazrin family.Contains 3 SAM (sterile alpha motif) domains
SUBUNIT	Isoform 2, isoform 3 and isoform 4 interact with PPLN-terminus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Component of the cornified envelope of keratinocytes. May be involved in the interplay between adherens junctions and desmosomes. The function in the nucleus is not known.

Kazrin Polyclonal Antibody - Additional Information

Gene ID 23254

Other Names

Kazrin, KAZN (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=29173 target="_blank">HGNC:29173)

Target/Specificity

Isoform 2, isoform 3 and isoform 4 are expressed in several cell lines including keratinocytes and bladder and epidermoid carcinoma (at protein level). Isoform 2, isoform 3 and isoform 4 are expressed in hair follicle and interfollicular epidermis (at protein level).

Dilution

WB~1:1000

="dilution_IHC-P">IHC-P~N/A<br \><span class
="dilution_IHC-F">IHC-F~N/A<br \><span class
="dilution_IF">IF~1:50~200<br \>E~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Kazrin Polyclonal Antibody - Protein Information

Name KAZN ([HGNC:29173](#))

Function

Component of the cornified envelope of keratinocytes. May be involved in the interplay between adherens junctions and desmosomes. The function in the nucleus is not known.

Cellular Location

Cytoplasm, cytoskeleton. [Isoform 3]: Cytoplasm. Cell junction, desmosome. Nucleus.
Note=Observed at the apical plasma membrane of keratinocytes Partially colocalizes with PPL and DP at desmosomes, and with PP at the interdesmosomal plasma membrane. Colocalizes with cortical actin-based membrane structures

Tissue Location

Isoform 2, isoform 3 and isoform 4 are expressed in several cell lines including keratinocytes and bladder and epidermoid carcinoma (at protein level). Isoform 2, isoform 3 and isoform 4 are expressed in hair follicle and interfollicular epidermis (at protein level).

Kazrin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Kazrin Polyclonal Antibody - Images