

C10orf27 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59359**Specification**

C10orf27 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q96M53
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39460

C10orf27 Polyclonal Antibody - Additional Information**Gene ID** 219793**Other Names**

Protein TBATA, Protein SPATIAL, Stromal protein associated with thymii and lymph node homolog, Thymus, brain and testes-associated protein, TBATA, C10orf27, SPATIAL

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C10orf27 Polyclonal Antibody - Protein Information**Name** TBATA**Synonyms** C10orf27, SPATIAL**Function**

May play a role in spermatid differentiation. Modulates thymic stromal cell proliferation and thymus function.

Cellular Location

Cytoplasm, cytosol.


C10orf27 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C10orf27 Polyclonal Antibody - Images



Sample: A549 Cell (Human) Lysate at 40 ug
Primary: Anti-C10orf27 (bs-9720R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 39 kD
Observed band size: 42 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C10orf27) Polyclonal Antibody, Unconjugated (bs-9720R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.