

BCL7C Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59377

Specification

BCL7C Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, E

Primary Accession <u>O8WUZ0</u>

Reactivity
Host
Clonality
Rat, Pig, Dog, Bovine
Rabbit
Polyclonal

Clonality Polyclo
Calculated MW 23 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human BCL7C

Epitope Specificity 9-100/217

Isotype IgG
Purity

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.
SIMILARITY Belongs to the BCL7 family.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

This gene is identified by the similarity of its product to the N-terminal region of BCL7A protein. The BCL7A protein is encoded by the gene known to be directly involved in a three-way gene translocation in a Burkitt lymphoma cell line. The function of this gene has not yet been determined. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2013]

BCL7C Polyclonal Antibody - Additional Information

Gene ID 9274

Other Names

B-cell CLL/lymphoma 7 protein family member C, BCL7C

Target/Specificity

Ubiquitous.

Dilution

WB~~1:1000<br \><span class</pre>

="dilution IHC-P">IHC-P~~N/A<br \><span class

="dilution IHC-F">IHC-F~~N/A<br \><span class

="dilution_IF">IF \sim 1:50 \sim 200<br\>E \sim N/A



Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

BCL7C Polyclonal Antibody - Protein Information

Name BCL7C

FunctionMay play an anti-apoptotic role.


Tissue Location Ubiquitous..

BCL7C Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

BCL7C Polyclonal Antibody - Images



Sample:

Siha (Mouse) Lysate at 40 ug

Primary: Anti- BCL7C (bs-9763R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 23 kD Observed band size: 23 kD

