

ECRG4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59388

Specification

ECRG4 Polyclonal Antibody - Product Information

Application Primary Accession

Reactivity
Host
Clonality
Calculated MW
Physical State
Immunogen

Epitope Specificity

Isotype **Purity**

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

from human ECRG4/C2orf40

WB, IHC-P, IHC-F, IF, E

KLH conjugated synthetic peptide derived

SUBCELLULAR LOCATION Secreted. Cytoplasmic vesicle, secretory

vesicle.

O9H1Z8

Rabbit

7.8 KDa

Liquid

laG

Polyclonal

41-148/148

Rat, Pig, Dog

SIMILARITY Belongs to the augurin family.

Important Note

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

ECRG4, also known as augurin or C2orf40, is a 148 amino acid secreted protein. Belonging to the augurin family, ECRG4 is thought to be a hormone. It has also been suggested that ECRG4 may act as a tumor suppressor. The gene that encodes ECRG4 maps to human chromosome 2, which consists of 237 million bases encoding over 1,400 genes, making up approximately 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin icthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alstr syndrome is due to mutations in the ALMS1 gene. Interestingly, chromosome 2 contains what appears to be a vestigial second centromere and vestigial telomeres which gives credence to the hypothesis that human chromosome 2 is the result of an ancient fusion of two ancestral chromosomes.

ECRG4 Polyclonal Antibody - Additional Information

Gene ID 84417

Other Names

Augurin, Esophageal cancer-related gene 4 protein {ECO:0000303|Ref.1}, ECRG4, ECRG4 (HGNC:24642), C2orf40



Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

ECRG4 Polyclonal Antibody - Protein Information

Name ECRG4 (HGNC:24642)

Synonyms C2orf40

Function

Probable hormone that may attenuate cell proliferation and induce senescence of oligodendrocyte and neural precursor cells in the central nervous system (By similarity). ECRG4-induced senescence is characterized by G1 arrest, RB1 dephosphorylation and accelerated CCND1 and CCND3 proteasomal degradation (By similarity).

Cellular Location

Secreted. Cytoplasm. Apical cell membrane

Tissue Location

Expressed in the brain, with expression in the epithelial cell layer of the choroid plexus (at protein level)

ECRG4 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ECRG4 Polyclonal Antibody - Images







Paraformaldehyde-fixed, paraffin embedded (Rat adrenal gland); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ECRG4) Polyclonal Antibody, Unconjugated (bs-9807R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.