

C2orf44 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59391**Specification**

C2orf44 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q9H6R7
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79136

C2orf44 Polyclonal Antibody - Additional Information**Gene ID** 80304**Other Names**

WD repeat and coiled-coil-containing protein, WDCP (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=26157), C2orf44

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


C2orf44 Polyclonal Antibody - Protein Information**Name** WDCP ([HGNC:26157](#))**Synonyms** C2orf44**C2orf44 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

C2orf44 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (liver cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C2orf44) Polyclonal Antibody, Unconjugated (bs-9814R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.