

Adropin Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59415**Specification****Adropin Polyclonal Antibody - Product Information**

| | |
|--------------------------------|---|
| Application | IHC-P, IHC-F, IF, E |
| Primary Accession | Q6UWT2 |
| Reactivity | Pig, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 5 KDa |
| Physical State | Liquid |
| Immunogen | KLH conjugated synthetic peptide derived from human Adropin |
| Epitope Specificity | 21-76/76 |
| Isotype | IgG |
| Purity | |
| affinity purified by Protein A | |
| Buffer | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| SUBCELLULAR LOCATION | Secreted. |
| Important Note | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |

Background Descriptions

Involved in the regulation of glucose homeostasis and lipid metabolism.

Adropin Polyclonal Antibody - Additional Information

Gene ID 375704

Other Names

Adropin, Energy homeostasis-associated protein, ENHO, C9orf165

Target/Specificity

Expressed in liver and brain.

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A
IF ~ ~ 1:50 ~ 200
E ~ ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH

7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Adropin Polyclonal Antibody - Protein Information

Name ENHO

Synonyms C9orf165

Function

Involved in the regulation of glucose homeostasis and lipid metabolism.

Cellular Location

Secreted.

Tissue Location


Expressed in liver and brain.

Adropin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Adropin Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Adropin) Polyclonal Antibody, Unconjugated (bs-9886R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.