

**Adropin Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP59415****Specification**

---

**Adropin Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">Q6UWT2</a>
Reactivity	Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	7927

**Adropin Polyclonal Antibody - Additional Information****Gene ID** 375704**Other Names**

Adropin, Energy homeostasis-associated protein, ENHO, C9orf165

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
E ~ ~ N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Adropin Polyclonal Antibody - Protein Information****Name** ENHO**Synonyms** C9orf165**Function**

Involved in the regulation of glucose homeostasis and lipid metabolism.

**Cellular Location**

Secreted.

**Tissue Location**


Expressed in liver and brain.

## **Adropin Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Adropin Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Adropin) Polyclonal Antibody, Unconjugated (bs-9886R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.