

C11ORF77 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59429**Specification**

C11ORF77 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q96MB7
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39146

C11ORF77 Polyclonal Antibody - Additional Information**Gene ID** 283254**Other Names**

Putative nuclease HARBI1, 3.1.-., Harbinger transposase-derived nuclease, HARBI1, C11orf77

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C11ORF77 Polyclonal Antibody - Protein Information**Name** HARBI1**Synonyms** C11orf77**Function**

Transposase-derived protein that may have nuclease activity (Potential). Does not have transposase activity.

Cellular Location

Nucleus. Cytoplasm. Note=Interaction with NAIF1 promotes translocation to the nucleus

Tissue Location


Detected in brain, eye, nerve tissue, kidney and lung.

C11ORF77 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C11ORF77 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C11ORF77) Polyclonal Antibody, Unconjugated (bs-9934R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.