

**CREG2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP59443****Specification**

---

**CREG2 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q8IUH2</a>
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32109

**CREG2 Polyclonal Antibody - Additional Information****Gene ID** 200407**Other Names**

Protein CREG2, Cellular repressor of E1A-stimulated genes 2, CREG2

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**CREG2 Polyclonal Antibody - Protein Information****Name** CREG2**Cellular Location**

Secreted.

**Tissue Location**

Brain specific mainly in the limbic system and faintly in the spinal cord but not in cerebellum


**CREG2 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **CREG2 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CREG2) Polyclonal Antibody, Unconjugated (bs-9957R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CREG2) Polyclonal Antibody, Unconjugated (bs-9957R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.