

C21orf70 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59452**Specification****C21orf70 Polyclonal Antibody - Product Information**

| | |
|---------------------|---|
| Application | WB, IHC-P, IHC-F, IF, ICC, E |
| Primary Accession | Q9NSI2 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 25 KDa |
| Physical State | Liquid |
| Immunogen | KLH conjugated synthetic peptide derived from human C21orf70 |
| Epitope Specificity | 51-150/230 |
| Isotype | IgG |
| Purity | affinity purified by Protein A |
| Buffer | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| Important Note | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |

C21orf70 Polyclonal Antibody - Additional Information**Gene ID** 85395**Other Names**

Protein FAM207A, FAM207A, C21orf70

Dilution

WB~~1:1000
IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
ICC~~N/A
E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C21orf70 Polyclonal Antibody - Protein Information**Name** SLX9 ([HGNC:15811](#))**Synonyms** C21orf70, FAM207A

Function

May be involved in ribosome biogenesis.

Cellular Location

Nucleus, nucleolus {ECO:0000250|UniProtKB:P53251}


Tissue Location

Not detected in any tested tissue.

C21orf70 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C21orf70 Polyclonal Antibody - Images

Paraformaldehyde-fixed, paraffin embedded (human stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C21orf70) Polyclonal Antibody, Unconjugated (bs-9979R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.