

C4orf22 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59461**Specification**

C4orf22 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q6V702
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26869

C4orf22 Polyclonal Antibody - Additional Information**Gene ID** 255119**Other Names**

Cilia- and flagella-associated protein 299, CFAP299 (HGNC:28554)

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C4orf22 Polyclonal Antibody - Protein Information**Name** CFAP299 ([HGNC:28554](#))**Function**

May be involved in spermatogenesis.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q810M1}. Nucleus {ECO:0000250|UniProtKB:Q810M1}.
Note=Mainly cytoplasmic {ECO:0000250|UniProtKB:Q810M1}


C4orf22 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C4orf22 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C4orf22) Polyclonal Antibody, Unconjugated (bs-9991R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.