

**Anti-ATF1 Antibody**  
**Rabbit polyclonal antibody to ATF1**  
**Catalog # AP59483**

### Specification

#### Anti-ATF1 Antibody - Product Information

Application	WB, IP, IF/IC, IHC
Primary Accession	<a href="#">P18846</a>
Other Accession	<a href="#">P81269</a>
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29233

#### Anti-ATF1 Antibody - Additional Information

##### Gene ID 466

##### Other Names

Cyclic AMP-dependent transcription factor ATF-1; cAMP-dependent transcription factor ATF-1; Activating transcription factor 1; Protein TREB36

##### Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human ATF1. The exact sequence is proprietary.

##### Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 - 1/100)  
IP~~N/A  
IF/IC~~N/A  
IHC~~1:100~500

##### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

##### Storage

Store at -20 °C. Stable for 12 months from date of receipt

#### Anti-ATF1 Antibody - Protein Information

##### Name ATF1

##### Function

This protein binds the cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Binds to the Tax-responsive element (TRE) of HTLV-I. Mediates PKA-induced stimulation of CRE-reporter genes. Represses the expression of FTH1 and other antioxidant detoxification genes. Triggers cell proliferation and transformation.

## Cellular Location


Nucleus.

## Anti-ATF1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## Anti-ATF1 Antibody - Images



Western blot analysis of ATF1 expression in HCT116 (A), HEK293T (B), mouse kidney (C), PC12 (D), Panc1 (E) whole cell lysates.



Immunohistochemical analysis of ATF1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ATF1 staining in KNRK cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

#### **Anti-ATF1 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human ATF1. The exact sequence is proprietary.