

**Anti-BST2 Antibody**  
**Rabbit polyclonal antibody to BST2**  
**Catalog # AP59491**

**Specification**

---

**Anti-BST2 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q10589</a>
Other Accession	<a href="#">Q8R2Q8</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19769

**Anti-BST2 Antibody - Additional Information**

**Gene ID** 684

**Other Names**

Bone marrow stromal antigen 2; BST-2; HM1.24 antigen; Tetherin; CD317

**Target/Specificity**

Recognizes endogenous levels of BST2 protein.

**Dilution**

WB~~WB (1/500 - 1/1000)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-BST2 Antibody - Protein Information**

**Name** BST2

**Function**

IFN-induced antiviral host restriction factor which efficiently blocks the release of diverse mammalian enveloped viruses by directly tethering nascent virions to the membranes of infected cells. Acts as a direct physical tether, holding virions to the cell membrane and linking virions to each other. The tethered virions can be internalized by endocytosis and subsequently degraded or they can remain on the cell surface. In either case, their spread as cell-free virions is restricted (PubMed:<a href="http://www.uniprot.org/citations/18200009" target="\_blank">18200009</a>, PubMed:<a href="http://www.uniprot.org/citations/18342597" target="\_blank">18342597</a>, PubMed:<a href="http://www.uniprot.org/citations/19036818" target="\_blank">19036818</a>, PubMed:<a href="http://www.uniprot.org/citations/19879838" target="\_blank">19879838</a>,

PubMed:<a href="http://www.uniprot.org/citations/20019814" target="\_blank">20019814</a>, PubMed:<a href="http://www.uniprot.org/citations/20399176" target="\_blank">20399176</a>, PubMed:<a href="http://www.uniprot.org/citations/20419159" target="\_blank">20419159</a>, PubMed:<a href="http://www.uniprot.org/citations/20940320" target="\_blank">20940320</a>, PubMed:<a href="http://www.uniprot.org/citations/21529378" target="\_blank">21529378</a>, PubMed:<a href="http://www.uniprot.org/citations/22520941" target="\_blank">22520941</a>, PubMed:<a href="http://www.uniprot.org/citations/37922253" target="\_blank">37922253</a>). Its target viruses belong to diverse families, including retroviridae: human immunodeficiency virus type 1 (HIV-1), human immunodeficiency virus type 2 (HIV-2), simian immunodeficiency viruses (SIVs), equine infectious anemia virus (EIAV), feline immunodeficiency virus (FIV), prototype foamy virus (PFV), Mason-Pfizer monkey virus (MPMV), human T-cell leukemia virus type 1 (HTLV-1), Rous sarcoma virus (RSV) and murine leukemia virus (MLV), flaviviridae: hepatitis C virus (HCV), filoviridae: ebola virus (EBOV) and marburg virus (MARV), arenaviridae: lassa virus (LASV) and machupo virus (MACV), herpesviridae: kaposi sarcoma-associated herpesvirus (KSHV), rhabdoviridae: vesicular stomatitis virus (VSV), orthomyxoviridae: influenza A virus, paramyxoviridae: nipah virus, and coronaviridae: SARS-CoV (PubMed:<a href="http://www.uniprot.org/citations/18200009" target="\_blank">18200009</a>, PubMed:<a href="http://www.uniprot.org/citations/18342597" target="\_blank">18342597</a>, PubMed:<a href="http://www.uniprot.org/citations/19179289" target="\_blank">19179289</a>, PubMed:<a href="http://www.uniprot.org/citations/19879838" target="\_blank">19879838</a>, PubMed:<a href="http://www.uniprot.org/citations/20399176" target="\_blank">20399176</a>, PubMed:<a href="http://www.uniprot.org/citations/20419159" target="\_blank">20419159</a>, PubMed:<a href="http://www.uniprot.org/citations/20686043" target="\_blank">20686043</a>, PubMed:<a href="http://www.uniprot.org/citations/20943977" target="\_blank">20943977</a>, PubMed:<a href="http://www.uniprot.org/citations/21529378" target="\_blank">21529378</a>, PubMed:<a href="http://www.uniprot.org/citations/21621240" target="\_blank">21621240</a>, PubMed:<a href="http://www.uniprot.org/citations/22520941" target="\_blank">22520941</a>, PubMed:<a href="http://www.uniprot.org/citations/26378163" target="\_blank">26378163</a>, PubMed:<a href="http://www.uniprot.org/citations/31199522" target="\_blank">31199522</a>). Can inhibit cell surface proteolytic activity of MMP14 causing decreased activation of MMP15 which results in inhibition of cell growth and migration (PubMed:<a href="http://www.uniprot.org/citations/22065321" target="\_blank">22065321</a>). Can stimulate signaling by LILRA4/ILT7 and consequently provide negative feedback to the production of IFN by plasmacytoid dendritic cells in response to viral infection (PubMed:<a href="http://www.uniprot.org/citations/19564354" target="\_blank">19564354</a>, PubMed:<a href="http://www.uniprot.org/citations/26172439" target="\_blank">26172439</a>). Plays a role in the organization of the subapical actin cytoskeleton in polarized epithelial cells. Isoform 1 and isoform 2 are both effective viral restriction factors but have differing antiviral and signaling activities (PubMed:<a href="http://www.uniprot.org/citations/23028328" target="\_blank">23028328</a>, PubMed:<a href="http://www.uniprot.org/citations/26172439" target="\_blank">26172439</a>). Isoform 2 is resistant to HIV-1 Vpu-mediated degradation and restricts HIV-1 viral budding in the presence of Vpu (PubMed:<a href="http://www.uniprot.org/citations/23028328" target="\_blank">23028328</a>, PubMed:<a href="http://www.uniprot.org/citations/26172439" target="\_blank">26172439</a>). Isoform 1 acts as an activator of NF-kappa-B and this activity is inhibited by isoform 2 (PubMed:<a href="http://www.uniprot.org/citations/23028328" target="\_blank">23028328</a>).

### Cellular Location

Golgi apparatus, trans-Golgi network. Cell membrane; Single-pass type II membrane protein. Cell membrane; Lipid- anchor, GPI-anchor. Membrane raft. Cytoplasm. Apical cell membrane. Note=Shuttles between the cell membrane, where it is present predominantly in membrane/lipid rafts, and the trans- Golgi network. Forms a complex with MMP14 and localizes to the cytoplasm

### Tissue Location

Predominantly expressed in liver, lung, heart and placenta. Lower levels in pancreas, kidney, skeletal muscle and brain Overexpressed in multiple myeloma cells. Highly expressed during B-cell development, from pro-B precursors to plasma cells. Highly expressed on T-cells, monocytes, NK


cells and dendritic cells (at protein level)

### Anti-BST2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-BST2 Antibody - Images



Western blot analysis of BST2 expression in HeLa (A), mouse spleen (B), mouse lung (C), rat spleen (D) whole cell lysates.

### Anti-BST2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human BST2. The exact sequence is proprietary.