

## **Anti-Calmodulin Antibody**

Rabbit polyclonal antibody to Calmodulin Catalog # AP59496

## **Specification**

## **Anti-Calmodulin Antibody - Product Information**

Application WB, IP, IF/IC, IHC

Primary Accession P62158
Other Accession P62204

Reactivity Human, Mouse, Rat, Rabbit, Zebrafish,

Chicken, Bovine, SARS

Host Rabbit Clonality Polyclonal

# **Anti-Calmodulin Antibody - Additional Information**

#### **Other Names**

CALM; CAM; CAM1; CAM2; CAMB; CALML2; CAM3; CAMC; CAMIII; Calmodulin; CaM

## Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Calmodulin. The exact sequence is proprietary.

#### **Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 - 1/100) IP~~N/A IF/IC~~N/A IHC~~1:100~500

#### **Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

### **Storage**

Store at -20 °C. Stable for 12 months from date of receipt

## **Anti-Calmodulin Antibody - Protein Information**

## **Anti-Calmodulin Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry




- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **Anti-Calmodulin Antibody - Images**




Western blot analysis of Calmodulin expression in HEK293T (A), Hela (B), mouse testis (C), mouse kidney (D), rat testis (E) whole cell lysates.



Immunohistochemical analysis of Calmodulin staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.





Immunofluorescent analysis of Calmodulin staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

## **Anti-Calmodulin Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Calmodulin. The exact sequence is proprietary.