

Anti-IRAK1 (pT209) Antibody
Rabbit polyclonal antibody to IRAK1 (pT209)
Catalog # AP59591

Specification

Anti-IRAK1 (pT209) Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P51617 |
| Other Accession | Q62406 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 76537 |

Anti-IRAK1 (pT209) Antibody - Additional Information

Gene ID 3654

Other Names

IRAK; Interleukin-1 receptor-associated kinase 1; IRAK-1

Target/Specificity

Recognizes endogenous levels of IRAK1 (pT209) protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-IRAK1 (pT209) Antibody - Protein Information

Name IRAK1 ([HGNC:6112](#))

Synonyms IRAK

Function

Serine/threonine-protein kinase that plays a critical role in initiating innate immune response against foreign pathogens. Involved in Toll-like receptor (TLR) and IL-1R signaling pathways. Is rapidly recruited by MYD88 to the receptor-signaling complex upon TLR activation. Association with MYD88 leads to IRAK1 phosphorylation by IRAK4 and subsequent autophosphorylation and kinase activation. Phosphorylates E3 ubiquitin ligases Pellino proteins (PELI1, PELI2 and PELI3) to promote pellino-mediated polyubiquitination of IRAK1. Then, the ubiquitin-binding domain of IKBKG/NEMO binds to polyubiquitinated IRAK1 bringing together the IRAK1-MAP3K7/TAK1-TRAF6

complex and the NEMO-IKKA-IKKB complex. In turn, MAP3K7/TAK1 activates IKKs (CHUK/IKKA and IKBKB/IKKB) leading to NF-kappa-B nuclear translocation and activation. Alternatively, phosphorylates TIRAP to promote its ubiquitination and subsequent degradation. Phosphorylates the interferon regulatory factor 7 (IRF7) to induce its activation and translocation to the nucleus, resulting in transcriptional activation of type I IFN genes, which drive the cell in an antiviral state. When sumoylated, translocates to the nucleus and phosphorylates STAT3.

Cellular Location

Cytoplasm. Nucleus. Lipid droplet Note=Translocates to the nucleus when sumoylated. RSAD2/viperin recruits it to the lipid droplet (By similarity).

Tissue Location


Isoform 1 and isoform 2 are ubiquitously expressed in all tissues examined, with isoform 1 being more strongly expressed than isoform 2.

Anti-IRAK1 (pT209) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IRAK1 (pT209) Antibody - Images



Western blot analysis of IRAK1 (pT209) expression in A2780 (A), rat thymus (B) whole cell lysates.

Anti-IRAK1 (pT209) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human IRAK1. The exact sequence is proprietary.