

Anti-HSL Antibody
Rabbit polyclonal antibody to HSL
Catalog # AP59608**Specification**

Anti-HSL Antibody - Product Information

Application	WB, IP, IHC
Primary Accession	Q05469
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	116598

Anti-HSL Antibody - Additional Information**Gene ID** 3991**Other Names**

Hormone-sensitive lipase; HSL

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HSL. The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IP (1/10 - 1/100)

IP~~N/A

IHC~~1:100~500

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-HSL Antibody - Protein Information**Name** LIPE**Function**

Lipase with broad substrate specificity, catalyzing the hydrolysis of triacylglycerols (TAGs), diacylglycerols (DAGs), monoacylglycerols (MAGs), cholesteryl esters and retinyl esters (PubMed: [15716583](http://www.uniprot.org/citations/15716583), PubMed: [15955102](http://www.uniprot.org/citations/15955102), PubMed: [19800417](http://www.uniprot.org/citations/19800417), PubMed: [8812477](http://www.uniprot.org/citations/8812477)). Shows a preferential hydrolysis of DAGs over TAGs and MAGs and preferentially hydrolyzes the

fatty acid (FA) esters at the sn-3 position of the glycerol backbone in DAGs (PubMed:19800417). Preferentially hydrolyzes FA esters at the sn-1 and sn-2 positions of the glycerol backbone in TAGs (By similarity). Catalyzes the hydrolysis of 2-arachidonoylglycerol, an endocannabinoid and of 2-acetyl monoalkylglycerol ether, the penultimate precursor of the pathway for de novo synthesis of platelet-activating factor (By similarity). In adipose tissue and heart, it primarily hydrolyzes stored triglycerides to free fatty acids, while in steroidogenic tissues, it principally converts cholesteryl esters to free cholesterol for steroid hormone production (By similarity).

Cellular Location

Cell membrane. Membrane, caveola. Cytoplasm, cytosol. Lipid droplet {ECO:0000250|UniProtKB:P54310}. Note=Found in the high-density caveolae. Translocates to the cytoplasm from the caveolae upon insulin stimulation (PubMed:17026959). Phosphorylation by AMPK reduces its translocation towards the lipid droplets (By similarity) {ECO:0000250|UniProtKB:P54310, ECO:0000269|PubMed:17026959}

Tissue Location


Testis..

Anti-HSL Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-HSL Antibody - Images



Western blot analysis of HSL expression in Panc1 (A), SKOVCA3 (B), C6 (C), 3T3L1 (D) whole cell lysates.



Immunohistochemical analysis of HSL staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-HSL Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HSL. The exact sequence is proprietary.