

Anti-NFYA Antibody

Rabbit polyclonal antibody to NFYA Catalog # AP59640

Specification

Anti-NFYA Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Calculated MW WB, IP, IF/IC, IHC <u>P23511</u> <u>P23708</u> Human, Mouse, Rat, Pig, Bovine, Dog Rabbit Polyclonal 36877

Anti-NFYA Antibody - Additional Information

Gene ID 4800

Other Names Nuclear transcription factor Y subunit alpha; CAAT box DNA-binding protein subunit A; Nuclear transcription factor Y subunit A; NF-YA

Target/Specificity Recognizes endogenous levels of NFYA protein.

Dilution WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 - 1/100) IP~~N/A IF/IC~~N/A IHC~~1:100~500

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Anti-NFYA Antibody - Protein Information

Name NFYA

Function

Component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'-CCAAT-3' box motif found in the promoters of its target genes. NF-Y can function as both an activator and a repressor, depending on its interacting cofactors. NF-YA positively regulates the transcription of the core clock component BMAL1.



Cellular Location Nucleus.

Anti-NFYA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-NFYA Antibody - Images



Western blot analysis of NFYA expression in Hela (A), mouse testis (B), rat lung (C) whole cell lysates.



Immunohistochemical analysis of NFYA staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.





Immunofluorescent analysis of NFYA staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Anti-NFYA Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human NFYA. The exact sequence is proprietary.