

Anti-OCRL Antibody
Rabbit polyclonal antibody to OCRL
Catalog # AP59647**Specification**

Anti-OCRL Antibody - Product Information

Application	WB
Primary Accession	Q01968
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	104205

Anti-OCRL Antibody - Additional Information**Gene ID** 4952**Other Names**

INPP5F; OCRL1; Inositol polyphosphate 5-phosphatase OCRL-1; Lowe oculocerebrorenal syndrome protein

Target/Specificity

Recognizes endogenous levels of OCRL protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-OCRL Antibody - Protein Information**Name** OCRL ([HGNC:8108](#))**Synonyms** OCRL1 {ECO:0000303|Ref.2}**Function**

Catalyzes the hydrolysis of the 5-position phosphate of phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P2) and phosphatidylinositol-3,4,5-bisphosphate (PtdIns(3,4,5)P3), with the greatest catalytic activity towards PtdIns(4,5)P2 (PubMed:10764818, PubMed:15474001, PubMed:7761412, PubMed:9430698). Able also to

hydrolyze the 5-phosphate of inositol 1,4,5-trisphosphate and of inositol 1,3,4,5-tetrakisphosphate (PubMed:25869668, PubMed:7761412). Regulates traffic in the endosomal pathway by regulating the specific pool of phosphatidylinositol 4,5-bisphosphate that is associated with endosomes (PubMed:21971085). Involved in primary cilia assembly (PubMed:22228094, PubMed:22543976). Acts as a regulator of phagocytosis, hydrolyzing PtdIns(4,5)P2 to promote phagosome closure, through attenuation of PI3K signaling (PubMed:22072788).

Cellular Location

Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:D3ZGS3}. Early endosome membrane. Membrane, clathrin-coated pit. Cell projection, cilium, photoreceptor outer segment. Cell projection, cilium. Cytoplasmic vesicle {ECO:0000250|UniProtKB:D3ZGS3}. Endosome. Golgi apparatus, trans-Golgi network {ECO:0000250|UniProtKB:D3ZGS3}. Lysosome Note=Also found on macropinosomes (PubMed:25869668). Colocalized with APPL1 on phagosomes (PubMed:22072788).

Tissue Location


Brain, skeletal muscle, heart, kidney, lung, placenta and fibroblasts. Expressed in the retina and the retinal pigment epithelium.

Anti-OCRL Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-OCRL Antibody - Images



Western blot analysis of OCRL expression in MG63 (A), A549 (B), HEK293T (C) whole cell lysates.

Anti-OCRL Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human OCRL. The exact sequence is proprietary.