

Anti-PAWR Antibody

Rabbit polyclonal antibody to PAWR Catalog # AP59653

Specification

Anti-PAWR Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Calculated MW WB, IHC <u>Q96IZ0</u> <u>Q925B0</u> Human, Mouse, Rat, Chicken, Bovine, Dog Rabbit Polyclonal 36568

Anti-PAWR Antibody - Additional Information

Gene ID 5074

Other Names PAR4; PRKC apoptosis WT1 regulator protein; Prostate apoptosis response 4 protein; Par-4

Target/Specificity Recognizes endogenous levels of PAWR protein.

Dilution WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200) IHC~~1:100~500

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Anti-PAWR Antibody - Protein Information

Name PAWR

Synonyms PAR4

Function

Pro-apoptotic protein capable of selectively inducing apoptosis in cancer cells, sensitizing the cells to diverse apoptotic stimuli and causing regression of tumors in animal models. Induces apoptosis in certain cancer cells by activation of the Fas prodeath pathway and coparallel inhibition of NF-kappa-B transcriptional activity. Inhibits the transcriptional activation and augments the transcriptional repression mediated by WT1. Down-regulates the anti- apoptotic protein BCL2 via its interaction with WT1. Also seems to be a transcriptional repressor by itself. May be directly



involved in regulating the amyloid precursor protein (APP) cleavage activity of BACE1.

Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic in absence of apoptosis signal and in normal cells. Nuclear in most cancer cell lines. Nuclear entry seems to be essential but not sufficient for apoptosis (By similarity). Nuclear localization includes nucleoplasm and PML nuclear bodies.

Tissue Location

Widely expressed. Expression is elevated in various neurodegenerative diseases such as amyotrophic lateral sclerosis, Alzheimer, Parkinson and Huntington diseases and stroke. Down-regulated in several cancers.

Anti-PAWR Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-PAWR Antibody - Images



Western blot analysis of PAWR expression in HEK293T (A), A549 (B), U2OS (C) whole cell lysates.





Immunohistochemical analysis of PAWR staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-PAWR Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PAWR. The exact sequence is proprietary.