

Specification

Anti-BCAS2 Antibody - Product Information

Anti-BCAS2 Antibody - Additional Information

Target/Specificity

Format

Anti-BCAS2 Antibody - Protein Information

[29360106](http://www.uniprot.org/citations/29360106), PubMed:<[30705154](http://www.uniprot.org/citations/30705154)>). Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing. May have a scaffolding role in the spliceosome assembly as it contacts all other components of the core complex. The PRP19-CDC5L complex may also play a role in the response to DNA damage (DDR).

Cellular Location

Nucleus. Nucleus, nucleolus


Tissue Location

Ubiquitously expressed.

Anti-BCAS2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-BCAS2 Antibody - Images

Western blot analysis of BCAS2 expression in HEK293T (A), Hela (B), U2OS (C), mouse kidney (D), mouse testis (E), rat kidney (F), rat testis (G) whole cell lysates.

Anti-BCAS2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human BCAS2. The exact sequence is proprietary.