

Anti-POLR3C Antibody
Rabbit polyclonal antibody to POLR3C
Catalog # AP59802**Specification**

Anti-POLR3C Antibody - Product Information

Application	WB, IP
Primary Accession	O9BUI4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60612

Anti-POLR3C Antibody - Additional Information**Gene ID** 10623**Other Names**

DNA-directed RNA polymerase III subunit RPC3; RNA polymerase III subunit C3; DNA-directed RNA polymerase III subunit C; RNA polymerase III 62 kDa subunit; RPC62

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human POLR3C. The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000), IP (1/10 - 1/100)

IP~~N/A

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-POLR3C Antibody - Protein Information**Name** POLR3C ([HGNC:30076](#))**Function**

DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates (PubMed:20413673, PubMed:33558764, PubMed:33558766, PubMed:34675218, PubMed:35637192). Specific

peripheric component of RNA polymerase III (Pol III) which synthesizes small non-coding RNAs including 5S rRNA, snRNAs, tRNAs and miRNAs from at least 500 distinct genomic loci (PubMed:20413673, PubMed:33558764, PubMed:33558766, PubMed:35637192). Part of POLR3C/RPC3-POLR3F/RPC6-POLR3G/RPC7 heterotrimer, coordinates the dynamics of Pol III stalk and clamp modules during the transition from apo to elongation state (PubMed:33558764, PubMed:33558766). Pol III plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as a nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs) induce type I interferon and NF-kappa-B through the RIG-I pathway (PubMed:19609254, PubMed:19631370). Preferentially binds single-stranded DNA (ssDNA) in a sequence-independent manner (PubMed:21358628).

Cellular Location


Nucleus.

Anti-POLR3C Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-POLR3C Antibody - Images



Western blot analysis of POLR3C expression in CT26 (A), PC12 (B), HeLa (C), COS7 (D), A549 (E) whole cell lysates.

Anti-POLR3C Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human POLR3C. The exact sequence is proprietary.