

**Anti-SERCA2 Antibody**  
**Rabbit polyclonal antibody to SERCA2**  
**Catalog # AP59916**

**Specification**

**Anti-SERCA2 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P16615</a>
Other Accession	<a href="#">055143</a>
Reactivity	Human, Mouse, Rat, Rabbit, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	114757

**Anti-SERCA2 Antibody - Additional Information**

**Gene ID 488**

**Other Names**

ATP2B; Sarcoplasmic/endoplasmic reticulum calcium ATPase 2; SERCA2; SR Ca(2+)-ATPase 2; Calcium pump 2; Calcium-transporting ATPase sarcoplasmic reticulum type, slow twitch skeletal muscle isoform; Endoplasmic reticulum class 1/2 Ca(2+) ATPase

**Target/Specificity**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SERCA2. The exact sequence is proprietary.

**Dilution**

WB~~WB (1/500 - 1/1000)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-SERCA2 Antibody - Protein Information**

**Name** ATP2A2 ([HGNC:812](#))

**Synonyms** ATP2B

**Function**

This magnesium-dependent enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen (PubMed:[12542527](http://www.uniprot.org/citations/12542527), PubMed:[16402920](http://www.uniprot.org/citations/16402920)). Involved in

autophagy in response to starvation. Upon interaction with VMP1 and activation, controls ER-isolation membrane contacts for autophagosome formation (PubMed:<a href="http://www.uniprot.org/citations/28890335" target="\_blank">28890335</a>). Also modulates ER contacts with lipid droplets, mitochondria and endosomes (PubMed:<a href="http://www.uniprot.org/citations/28890335" target="\_blank">28890335</a>). In coordination with FLVCR2 mediates heme-stimulated switching from mitochondrial ATP synthesis to thermogenesis (By similarity).

#### Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:O55143}; Multi-pass membrane protein. Sarcoplasmic reticulum membrane; Multi-pass membrane protein. Note=Colocalizes with FLVCR2 at the mitochondrial-ER contact junction. {ECO:0000250|UniProtKB:O55143}

#### Tissue Location


Isoform 1 is widely expressed in smooth muscle and nonmuscle tissues such as in adult skin epidermis, with highest expression in liver, pancreas and lung, and intermediate expression in brain, kidney and placenta. Also expressed at lower levels in heart and skeletal muscle. Isoforms 2 and 3 are highly expressed in the heart and slow twitch skeletal muscle. Expression of isoform 3 is predominantly restricted to cardiomyocytes and in close proximity to the sarcolemma. Both isoforms are mildly expressed in lung, kidney, liver, pancreas and placenta. Expression of isoform 3 is amplified during monocytic differentiation and also observed in the fetal heart.

#### Anti-SERCA2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-SERCA2 Antibody - Images



Western blot analysis of SERCA2 expression in THP1 (A), NIH3T3 (B), PC12 (C) whole cell lysates.

#### Anti-SERCA2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SERCA2. The exact sequence is proprietary.