

Anti-TRPV1 Antibody
Rabbit polyclonal antibody to TRPV1
Catalog # AP59942**Specification**

Anti-TRPV1 Antibody - Product Information

Application	WB
Primary Accession	Q8NER1
Other Accession	Q704Y3
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	94956

Anti-TRPV1 Antibody - Additional Information**Gene ID** 7442**Other Names**

VR1; Transient receptor potential cation channel subfamily V member 1; TrpV1; Capsaicin receptor; Osm-9-like TRP channel 1; OTRPC1; Vanilloid receptor 1

Target/Specificity

Recognizes endogenous levels of TRPV1 protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-TRPV1 Antibody - Protein Information**Name** TRPV1**Synonyms** VR1**Function**

Ligand-activated non-selective calcium permeant cation channel involved in detection of noxious chemical and thermal stimuli. Seems to mediate proton influx and may be involved in intracellular acidosis in nociceptive neurons. Involved in mediation of inflammatory pain and hyperalgesia. Sensitized by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases, which involves PKC isozymes and PCL. Activation by vanilloids, like capsaicin, and temperatures higher than 42 degrees Celsius, exhibits a time- and Ca(2+)-dependent outward

rectification, followed by a long-lasting refractory state. Mild extracellular acidic pH (6.5) potentiates channel activation by noxious heat and vanilloids, whereas acidic conditions (pH <6) directly activate the channel. Can be activated by endogenous compounds, including 12-hydroperoxytetraenoic acid and bradykinin. Acts as ionotropic endocannabinoid receptor with central neuromodulatory effects. Triggers a form of long-term depression (TRPV1-LTD) mediated by the endocannabinoid anandamine in the hippocampus and nucleus accumbens by affecting AMPA receptors endocytosis.

Cellular Location

Postsynaptic cell membrane {ECO:0000250|UniProtKB:O35433}; Multi-pass membrane protein {ECO:0000250|UniProtKB:O35433}. Cell projection, dendritic spine membrane {ECO:0000250|UniProtKB:O35433}; Multi-pass membrane protein {ECO:0000250|UniProtKB:O35433}. Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:O35433}. Note=Mostly, but not exclusively expressed in postsynaptic dendritic spines {ECO:0000250|UniProtKB:O35433}

Tissue Location


Widely expressed at low levels. Expression is elevated in dorsal root ganglia. In skin, expressed in cutaneous sensory nerve fibers, mast cells, epidermal keratinocytes, dermal blood vessels, the inner root sheet and the infundibulum of hair follicles, differentiated sebocytes, sweat gland ducts, and the secretory portion of eccrine sweat glands (at protein level)

Anti-TRPV1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TRPV1 Antibody - Images



Western blot analysis of TRPV1 expression in U87MG (A), A375 (B) whole cell lysates.

Anti-TRPV1 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human TRPV1. The exact sequence is proprietary.