

OTUD4 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5997a**Specification**

OTUD4 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q01804
Other Accession	NP_001096123.1 , NP_059963.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	124045
Antigen Region	403-430

OTUD4 Antibody (N-term) - Additional Information**Gene ID** 54726**Other Names**

OTU domain-containing protein 4, HIV-1-induced protein HIN-1, OTUD4, HIN1, KIAA1046

Target/Specificity

This OTUD4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 403-430 amino acids from the N-terminal region of human OTUD4.

Dilution

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OTUD4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OTUD4 Antibody (N-term) - Protein Information**Name** OTUD4 ([HGNC:24949](#))

Function Deubiquitinase which hydrolyzes the isopeptide bond between the ubiquitin C-terminus and the lysine epsilon-amino group of the target protein (PubMed:[23827681](#), PubMed:[25944111](#), PubMed:[29395066](#)). May negatively regulate inflammatory and pathogen recognition signaling in innate immune response. Upon phosphorylation at Ser-202 and Ser-204 residues, via IL-1 receptor and Toll-like receptor signaling pathway, specifically deubiquitinates 'Lys-63'-polyubiquitinated MYD88 adapter protein triggering down-regulation of NF-kappa-B-dependent transcription of inflammatory mediators (PubMed:[29395066](#)). Independently of the catalytic activity, acts as a scaffold for alternative deubiquitinases to assemble specific deubiquitinase- substrate complexes. Associates with USP7 and USP9X deubiquitinases to stabilize alkylation repair enzyme ALKBH3, thereby promoting the repair of alkylated DNA lesions (PubMed:[25944111](#)).


Cellular Location

Cytoplasm. Nucleus. Note=Primarily cytoplasmic


OTUD4 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OTUD4 Antibody (N-term) - Images

OTUD4 Antibody (N-term) (Cat. #AP5997a) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the OTUD4 antibody detected the OTUD4 protein (arrow).



OTUD4 antibody (N-term) (Cat. #AP5997a) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the OTUD4 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.