

Anti-hNRNP C Antibody
Rabbit polyclonal antibody to hNRNP C
Catalog # AP60015**Specification**

Anti-hNRNP C Antibody - Product Information

| | |
|-------------------|---|
| Application | WB, IF/IC |
| Primary Accession | P07910 |
| Reactivity | Human, Rabbit, Monkey, Pig, Bovine, Dog |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 33670 |

Anti-hNRNP C Antibody - Additional Information**Gene ID** 3183**Other Names**

HNRPC; Heterogeneous nuclear ribonucleoproteins C1/C2; hnRNP C1/C2

Target/Specificity

Recognizes endogenous levels of hNRNP C protein.

Dilution

WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

IF/IC~~N/A

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-hNRNP C Antibody - Protein Information**Name** HNRNPC**Synonyms** HNRPC**Function**

Binds pre-mRNA and nucleates the assembly of 40S hnRNP particles (PubMed: [8264621](http://www.uniprot.org/citations/8264621)). Interacts with poly-U tracts in the 3'-UTR or 5'-UTR of mRNA and modulates the stability and the level of translation of bound mRNA molecules (PubMed: [12509468](http://www.uniprot.org/citations/12509468), PubMed: [16010978](http://www.uniprot.org/citations/16010978), PubMed: [7567451](http://www.uniprot.org/citations/7567451), PubMed: [7567451](http://www.uniprot.org/citations/7567451)).

href="http://www.uniprot.org/citations/8264621" target="_blank">8264621). Single HNRNPC tetramers bind 230-240 nucleotides. Trimers of HNRNPC tetramers bind 700 nucleotides (PubMed:8264621). May play a role in the early steps of spliceosome assembly and pre-mRNA splicing. N6-methyladenosine (m6A) has been shown to alter the local structure in mRNAs and long non-coding RNAs (lncRNAs) via a mechanism named 'm(6)A-switch', facilitating binding of HNRNPC, leading to regulation of mRNA splicing (PubMed:25719671).

Cellular Location


Nucleus. Note=Component of ribonucleosomes

Anti-hNRNP C Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-hNRNP C Antibody - Images



Western blot analysis of hNRNP C expression in HeLa (A) whole cell lysates.



Immunofluorescent analysis of hNRNP C staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Anti-hNRNP C Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human hNRNP C. The exact sequence is proprietary.